

# Solutions

## 1.1. Estimates

Remark: your result might vary by some orders of magnitude from the ones below, that's OK!

1.) a fist:

$$L \sim \frac{G}{5c^5} \cdot \frac{(1 \text{ kg})^2 (0.3 \text{ m})^4}{(0.25 \text{ s})^6} \text{ with forearm}$$

$$\sim 10^{-51} \text{ W} \approx 10^{-51} \frac{1}{(0.25)^2} \cdot \frac{1}{5} \text{ gravitons}$$

$\sim 10^{-19}$  gravitons per second  
 $\approx 1$  graviton in 100 billion years ☹  
 cf. age of universe: 10 11

2.) steel rod:

$$L \sim \frac{G}{5c^5} \cdot \frac{(5 \cdot 10^5 \text{ kg})^2 (20 \text{ m})^4}{(0.1 \text{ s})^6} \sim 10^{-31} \text{ W} \quad \Downarrow \text{ still super small } \text{ ☹}$$

↑ men moves through system twice per period

3.) Earth:

$$L \sim \frac{G}{5c^5} \cdot \frac{(6 \cdot 10^{24} \text{ kg})^2 (3 \cdot 10^{11} \text{ m})^4}{(2 \cdot 10^7 \text{ s})^6}$$

just Earth, sun not moving  
 size of system  $\sim 2 \text{ AU}$   
 $\leftarrow \frac{1}{2} \text{ year}$

$\sim 0.1 \text{ W}$  a human produces  $\sim 100 \text{ W}$  of heat ☹

4.) black holes:

$$L \sim \frac{G}{5c^5} \cdot \frac{(10^{32} \text{ kg})^2 (5 \cdot 10^5 \text{ m})^4}{(0.015 \text{ s})^6}$$

both black holes  
 $r = \frac{G}{c^2} \cdot 6 \cdot 60 M_{\odot}$   
 $\leftarrow$  from Kepler's law  $T = 2\pi \sqrt{\frac{r^3}{G \cdot 60 M_{\odot}}} \sim 0.025$

$$\sim 10^{45} \text{ W} \approx \frac{0.01 M_{\odot} c^2}{5} \quad \Downarrow \text{ huge } \text{ ☹}$$

huge power, but still tiny amplitude/strain!

at 1AU:

$$\ddot{Q} \sim \sqrt{\frac{5c^5}{G}} \cdot L \sim 10^{49} \frac{\text{kg} \cdot \text{m}^2}{\text{s}^3}$$

at 10M:

$$h \sim \frac{G \cdot 2}{c^4} \cdot (0.015) \cdot \ddot{Q} \cdot \frac{1}{(10^{11} \text{ m})} \sim 10^{-8} \approx \frac{\text{hair}}{10 \text{ km}} \quad \Downarrow \text{ very small}$$

$$h \sim \frac{G \cdot 2}{c^4} \cdot (0.015) \cdot \ddot{Q} \cdot \frac{1}{10^{23} \text{ m}} \sim 10^{-20} \approx \frac{\text{proton}}{100 \text{ km}} \rightarrow \text{measured by LIGO } \text{ ☹}$$

1.2. binary system (circular)

with  $\phi = \omega t$ :  $\dot{\phi} = \omega$

$\dot{M}^{11} = -2\mu r^2 \omega \sin\phi \cos\phi$ ,  $\ddot{M}^{11} = -2\mu r^2 \omega^2 (\cos^2\phi - \sin^2\phi)$ ,  $\dddot{M}^{11} = \mu r^2 8\omega^3 \sin\phi \cos\phi$

$\dot{M}^{22} = 2\mu r^2 \omega \sin\phi \cos\phi$ ,  $\ddot{M}^{22} = -\ddot{M}^{11}$ ,  $\dddot{M}^{22} = -\dddot{M}^{11}$

$\dot{M}^{12} = \mu r^2 \omega (\cos^2\phi - \sin^2\phi)$ ,  $\ddot{M}^{12} = -\mu r^2 4\omega^2 \sin\phi \cos\phi$ ,  $\dddot{M}^{12} = -\mu r^2 4\omega^3 (2\cos^2\phi - 1)$

Luminosity:

$L = +\frac{1}{5} \langle \ddot{Q}_{ij} \ddot{Q}^{ij} \rangle$

$= \frac{1}{5} \langle \ddot{Q}_{ij} (\ddot{M}^{ij} - \frac{1}{3} \delta^{ij} \ddot{M}^k_k) \rangle$   
 $\rightarrow$  da  $\ddot{Q}_{ij} \delta^{ij} = 0$

$= \frac{1}{5} \langle \ddot{M}_{ij} \ddot{M}^{ij} - \frac{1}{3} (\ddot{M}^k_k)^2 \rangle$   
 $\ddot{M}^{11} + \ddot{M}^{22} = 0$

$(\ddot{M}^{11})^2 + 2(\ddot{M}^{12})^2 + (\ddot{M}^{22})^2$

$= \frac{1}{5} \int_{\text{orbit}} \frac{dt}{T} \cdot (4\mu r^2 \omega^3)^2 \cdot (8\sin^2\phi \cos^2\phi + 2(2\cos^2\phi - 1)^2)$   
 $\int_0^{2\pi} \frac{d\phi}{2\pi}$   
 $8\cos^2\phi(1 - \sin^2\phi) + 8\cos^4\phi - 8\cos^2\phi + 2 = 2$

$= \frac{1}{5} (4\mu r^2 \omega^3)^2 \cdot 2 \cdot \int_0^{2\pi} \frac{d\phi}{2\pi} \rightarrow 1$   
 $= \frac{1}{10} \mu^2 r^4 (2\omega)^6$

Using:  $r = (M\omega^{-2})^{1/3}$

$L = \frac{32}{5} \mu^2 M^{4/3} \omega^{10/3}$

$= \frac{32}{5} \underbrace{(\mu^{3/5} M^{2/5} \omega)}_{M_c}^{10/3}$

compare 1.7

3. Earth:

$M_c = 10^{27} \text{ kg}$ ,  $\omega = \frac{2\pi}{10^{7.5}} \text{ year}$

$L \approx 200 \text{ W}$

4. black holes:

$M_c = 5 \cdot 10^{31} \text{ kg}$ ,  $\omega = \sqrt{\frac{M}{r^3}} = 230 \text{ Hz}$

$L \approx 2 \cdot 10^{48} \text{ W} \approx 10 \frac{M_\odot}{s}$

WOW!

2.1. Wave equation

Christoffel:

$$2\Gamma_{\alpha\beta}^{\mu} = g^{\mu\nu} (\partial_{\alpha} g_{\beta\nu} + \partial_{\beta} g_{\alpha\nu} - \partial_{\nu} g_{\alpha\beta}), \quad g_{\alpha\beta} = \eta_{\alpha\beta} + h_{\alpha\beta}$$

$$\partial_{\alpha} g_{\beta\nu} = \partial_{\alpha} h_{\beta\nu} = \mathcal{O}(h)$$

$$= (\eta^{\mu\nu} + \mathcal{O}(h)) (\partial_{\alpha} h_{\beta\nu} + \partial_{\beta} h_{\alpha\nu} - \partial_{\nu} h_{\alpha\beta}) \quad \square$$

CURVATURE:

$$2R_{\mu\nu\alpha\beta} = 2(g_{\mu\rho} \partial_{\alpha} \Gamma_{\beta\nu}^{\rho} - g_{\mu\sigma} \partial_{\beta} \Gamma_{\alpha\nu}^{\sigma} + \underbrace{\Gamma_{\alpha\mu}^{\rho} \Gamma_{\beta\nu}^{\sigma} - \Gamma_{\beta\nu}^{\rho} \Gamma_{\alpha\mu}^{\sigma}}_{\mathcal{O}(h^2)})$$

$\uparrow \quad \uparrow$   
 $\eta + \mathcal{O}(h) \quad \eta + \mathcal{O}(h)$

$$= \partial_{\alpha} \partial_{\beta} h_{\nu\mu} + \partial_{\alpha} \partial_{\nu} h_{\beta\mu} - \partial_{\alpha} \partial_{\mu} h_{\beta\nu} - (\alpha \leftrightarrow \beta) + \mathcal{O}(h^2)$$

symmetric in  $\alpha, \beta \rightarrow 0$

$$= \partial_{\alpha} \partial_{\nu} h_{\beta\mu} + \partial_{\beta} \partial_{\mu} h_{\alpha\nu} - \partial_{\alpha} \partial_{\mu} h_{\beta\nu} - \partial_{\beta} \partial_{\nu} h_{\alpha\mu} + \mathcal{O}(h^2) \quad \square$$

$$2R_{\mu\nu} = 2R^{\alpha}{}_{\mu\alpha\nu} = \partial^{\alpha} \partial_{\mu} h_{\nu\alpha} + \partial^{\alpha} \partial^{\alpha} h_{\alpha\mu} - \partial^{\alpha} \partial_{\alpha} h_{\mu\nu} - \partial_{\mu} \partial_{\nu} \bar{h}^{\alpha}{}_{\alpha} + \mathcal{O}(h^2)$$

$h_{\mu\nu} = \bar{h}_{\mu\nu} - \eta_{\mu\nu} \bar{h}^{\alpha}{}_{\alpha} / 2$

$$= \partial^{\alpha} \partial_{\mu} \bar{h}_{\nu\alpha} - \partial_{\nu} \partial_{\mu} \bar{h}^{\alpha}{}_{\alpha} + \partial^{\alpha} \partial_{\nu} \bar{h}_{\mu\alpha} - \frac{1}{2} \partial^{\alpha} \partial_{\nu} \bar{h}^{\alpha}{}_{\alpha} - \square \bar{h}_{\mu\nu} + \frac{1}{2} \eta_{\mu\nu} \square \bar{h} + \partial_{\mu} \partial_{\nu} \bar{h}^{\alpha}{}_{\alpha}$$

$$= -\square \bar{h}_{\mu\nu} + \partial^{\alpha} \partial_{\mu} \bar{h}_{\nu\alpha} + \partial^{\alpha} \partial_{\nu} \bar{h}_{\mu\alpha} + \frac{1}{2} \eta_{\mu\nu} \square \bar{h}^{\alpha}{}_{\alpha} + \mathcal{O}(h^2) \quad (*)$$

$$2R = 2R^{\mu}{}_{\mu} = -\square \bar{h}^{\mu}{}_{\mu} + \partial^{\alpha} \partial^{\mu} \bar{h}_{\mu\alpha} + \partial^{\alpha} \partial^{\mu} \bar{h}_{\mu\alpha} + \frac{1}{2} \cdot 4 \cdot \square \bar{h}^{\alpha}{}_{\alpha} + \mathcal{O}(h^2)$$

$$= 2 \partial^{\alpha} \partial^{\beta} \bar{h}_{\alpha\beta} + \square \bar{h}^{\alpha}{}_{\alpha} \quad (**)$$

Einstein equations:

$$-2(R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R) \approx -(*) + \frac{1}{2} \eta_{\mu\nu} (**)$$

$$= \square \bar{h}_{\mu\nu} - \partial^{\alpha} \partial_{\mu} h_{\nu\alpha} - \partial^{\alpha} \partial_{\nu} h_{\mu\alpha} + \eta_{\mu\nu} \partial^{\alpha} \partial^{\beta} \bar{h}_{\alpha\beta} = -16\pi T_{\mu\nu} \quad \square$$

## 2.2 Gauge Trasfos

Ex P4

$$g'_{\mu\nu} = g_{\alpha\beta} \frac{\partial x^\alpha}{\partial x'^\mu} \frac{\partial x^\beta}{\partial x'^\nu}, \quad \frac{\partial x^\alpha}{\partial x'^\mu} = \delta_\mu^\alpha + \frac{\partial \xi^\alpha}{\partial x'^\mu} = \delta_\mu^\alpha + \frac{\partial \xi^\alpha}{\partial x^\mu} + \mathcal{O}(\eta^2) \approx \delta_\mu^\alpha + \partial_\mu \xi^\alpha$$

$x = x' + \mathcal{O}(\eta), \quad \xi = \mathcal{O}(\eta), \quad \partial_\mu = \partial'_\mu + \mathcal{O}(\eta)$

$$\eta_{\mu\nu} + h'_{\mu\nu} = (\eta_{\alpha\beta} + h_{\alpha\beta}) \left( \delta_\mu^\alpha + \partial_\mu \xi^\alpha \right) \left( \delta_\nu^\beta + \partial_\nu \xi^\beta \right)$$

$$= \underbrace{\eta_{\alpha\beta} \delta_\mu^\alpha \delta_\nu^\beta}_{\eta_{\mu\nu}} + \underbrace{h_{\alpha\beta} \delta_\mu^\alpha \delta_\nu^\beta}_{h_{\mu\nu}} + \underbrace{\eta_{\alpha\beta} \partial_\mu \xi^{\alpha\nu}}_{\partial_\mu \xi^\nu} + \underbrace{\eta_{\alpha\beta} \partial_\nu \xi^{\alpha\mu}}_{\partial_\nu \xi^\mu} + \mathcal{O}(\eta^2)$$

$$\hookrightarrow h'_{\mu\nu} = h_{\mu\nu} + \partial_\mu \xi_\nu + \partial_\nu \xi_\mu + \mathcal{O}(\eta^2)$$

for  $\bar{h}_{\mu\nu}$ :  $\bar{h}_{\mu\nu} = \bar{h}_{\mu\nu} - \frac{1}{2} \eta_{\mu\nu} \bar{h}^\alpha{}_\alpha$

$$\bar{h}'_{\mu\nu} = \bar{h}_{\mu\nu} - \frac{1}{2} \eta_{\mu\nu} \bar{h}'^\alpha{}_\alpha$$

$$= \bar{h}_{\mu\nu} + \partial_\mu \xi_\nu + \partial_\nu \xi_\mu - \frac{1}{2} \eta_{\mu\nu} (\bar{h}^\alpha{}_\alpha + 2 \partial_\alpha \xi^\alpha) + \mathcal{O}(\eta^2)$$

$$= \bar{h}_{\mu\nu} + \partial_\mu \xi_\nu + \partial_\nu \xi_\mu - \eta_{\mu\nu} \partial_\alpha \xi^\alpha + \mathcal{O}(\eta^2)$$

Now  $R'_{\mu\nu\alpha\beta}$ :

$$\partial_\nu \partial_\alpha h'_{\mu\beta} = \partial_\nu \partial_\alpha h_{\mu\beta} + \partial_\nu \partial_\alpha \partial_\mu \xi_\beta + \partial_\nu \partial_\alpha \partial_\beta \xi_\mu + \mathcal{O}(\eta^2)$$

$$\partial_\mu \partial_\beta h'_{\nu\alpha} = \partial_\mu \partial_\beta h_{\nu\alpha} + \partial_\mu \partial_\beta \partial_\nu \xi_\alpha + \partial_\mu \partial_\beta \partial_\alpha \xi_\nu + \mathcal{O}(\eta^2)$$

$$- \partial_\mu \partial_\alpha h'_{\nu\beta} = - \partial_\mu \partial_\alpha h_{\nu\beta} - \partial_\mu \partial_\alpha \partial_\nu \xi_\beta - \partial_\mu \partial_\alpha \partial_\beta \xi_\nu + \mathcal{O}(\eta^2)$$

$$- \partial_\nu \partial_\beta h'_{\mu\alpha} = - \partial_\nu \partial_\beta h_{\mu\alpha} - \partial_\nu \partial_\beta \partial_\mu \xi_\alpha - \partial_\nu \partial_\beta \partial_\alpha \xi_\mu + \mathcal{O}(\eta^2)$$

$\hookrightarrow$  all term with  $\xi$  cancel  $\square$

$$\hookrightarrow R'_{\mu\nu\alpha\beta} = R_{\mu\nu\alpha\beta} + \mathcal{O}(\eta^2)$$

harmonic gauge:

$$0 \stackrel{!}{=} \partial^\nu \bar{h}'_{\mu\nu} = \underbrace{\partial^\nu \bar{h}_{\mu\nu}}_{f_\mu} + \underbrace{\partial_\mu \partial^\nu \xi_\nu}_{\square} + \underbrace{\partial^\nu \partial_\nu \xi_\mu}_{-\partial_\mu \partial^\nu \xi_\nu} - \eta_{\mu\nu} \partial^\nu \partial_\alpha \xi^\alpha$$

$\hookrightarrow \square \xi_\mu = -f_\mu \hookrightarrow$  inhom. wave equation for  $\xi_\mu$  can (usually) be solved!

3.1. TT-gauge

1.)  $\bar{h}_{\mu\nu} = a_{\mu\nu} e^{ik_\alpha x^\alpha} + c.c.$  then:  $\partial_\mu \rightarrow ik_\mu$

$$\square \bar{h}_{\mu\nu} = 0 \quad \sim \quad \underbrace{k_\alpha k^\alpha}_{-k^0 k^0 + k^i k^i} = 0 \quad \sim \quad \omega^2 = (k^0)^2 = k^i k^i \quad (*)$$

gauge condition:  $\partial^\nu \bar{h}_{\mu\nu} = 0 \quad \sim \quad a_{\mu\nu} k^\nu = 0 \quad \checkmark$

2.)  $\square h_{\mu\nu} = \underbrace{\bar{h}_{\mu\nu}}_0 + \underbrace{\partial_\mu \square \partial_\nu}_0 + \underbrace{\partial_\nu \square \partial_\mu}_0 - \underbrace{\eta_{\mu\nu} \square \partial_\alpha \partial^\alpha \phi}_0 = 0 \quad \checkmark$

$$\partial^\nu h_{\mu\nu} = \underbrace{\partial^\nu \bar{h}_{\mu\nu}}_0 + \underbrace{\partial_\mu \partial^\nu \phi}_{-\square \phi_\mu = 0} + \underbrace{\partial_\nu \partial^\mu \phi}_{-\partial_\mu \partial^\nu \phi_{\nu\rho}} - \underbrace{\eta_{\mu\nu} \partial^\nu \partial^\alpha \phi}_0 = 0 \quad \checkmark$$

from (17):

$$a_{\mu\nu} e^{ik \cdot x} + c.c. = \left( a_{\mu\nu} + ik_\mu \frac{1}{i\omega} b_\nu + ik_\nu \frac{1}{i\omega} b_\mu - \eta_{\mu\nu} ik_\alpha \frac{1}{i\omega} b^\alpha \right) e^{ik \cdot x} + c.c.$$

$$\sim a_{\mu\nu} = a_{\mu\nu} + \eta_{\mu\nu} b_\nu + \eta_{\nu\mu} b_\mu - \eta_{\mu\nu} n_\alpha b^\alpha \quad \checkmark$$

with  $n_\mu = k_\mu / \omega$ ,  $n_0 = \frac{k_0}{\omega} = -1$ ,  $n_i n_i = \frac{k_i k_i}{\omega^2} = 1$   
 $k_0 = -\omega$

3.)  $a_{\mu}^{\mu} = 0$  &  $a_{0i} = 0$  ~~4 equations~~ in (19)

↳ 4 <sup>linear</sup> equations, can be solved for 4 components of  $b_\mu$   $\checkmark$

from (20):  $a_{\mu\nu} n^\nu = 0$

$\mu=0 \rightarrow a_{0\nu} n^\nu = 0 \sim a_{00} n^0 + a_{0j} n^j = 0 \sim a_{00} = 0 \quad \checkmark$

$\mu=i \rightarrow a_{i\nu} n^\nu = 0 \sim a_{i0} n^0 + a_{ij} n^j = 0 \sim a_{ij} n^j = 0 \quad \checkmark$

4.) ~~(22) is valid~~ <sup>correct</sup> if  $\square h_{ij}^{\pi} = 0$  and (21) is fulfilled

$$\square h_{ij}^{\pi} = \underbrace{\Lambda_{ij}^{kl}}_0 \underbrace{\square \bar{h}_{kl}}_0 = 0 \quad \checkmark$$

$$n^i \Lambda_{ijk} = \underbrace{n^i s_{ik}}_{n_k} - \underbrace{n^i n_i n_k}_1 = 0 \quad \sim \quad n^i \Lambda_{ij}^{kl} = 0 \quad \sim \quad n^i h_{ij}^{\pi} = 0 \quad \checkmark$$

$$s_{ij}^{\pi} \Lambda_{ij} = \underbrace{s_{ij}^{\pi} s_{ij}}_3 - \underbrace{s_{ij}^{\pi} n_i n_j}_1 = 2 \quad \sim \quad s_{ij}^{\pi} \Lambda_{ij}^{kl} = \Lambda_{ik}^{\pi} \Lambda_j^l - \underbrace{s_{ij}^{\pi} \Lambda_{ij}^{kl}}_2 \cdot \frac{1}{2} = 0$$

$$\sim h_i^{\pi\pi} = h_{ix}^{\pi\alpha} = 0 \quad \checkmark$$

3.1.4.) optional part: <sup>(alternative)</sup>

$$0 = a_{\mu}^{\pi\mu} = a_{\mu}^{\mu} + n^{\mu} b_{\mu} + n^{\mu} b_{\mu} - 4 \underbrace{n^{\mu} b^{\alpha}}_{n^{\mu} b_{\mu}} = a_{\mu}^{\mu} - 2 \underbrace{n^{\mu} b_{\mu}}_{n^0 b_0 + n^i b_i}$$

$$\Rightarrow b_0 = \frac{1}{2} a_{\mu}^{\mu} - n^i b_i$$

$$0 = a_{oi}^{\pi\pi} = a_{oi} + n_o b_i + n_i b_o \quad (*)$$

$$\hookrightarrow 0 = n^i ( \quad ) = a_{oi} n^i - b_i n^i + \frac{1}{2} a_{\mu}^{\mu} - \underbrace{n^j b_j}_{b_j n^j}$$

$$\Rightarrow b_i n^i = \frac{1}{2} a_{oi} n^i + \frac{1}{4} a_{\mu}^{\mu}$$

$$\Rightarrow b_0 = \frac{1}{4} a_{\mu}^{\mu} - \frac{1}{2} a_{oi} n^i$$

in (\*):  $0 = a_{oi} - b_i + n_i (\frac{1}{4} a_{\mu}^{\mu} - \frac{1}{2} a_{oj} n^j)$

$$\Rightarrow b_i = a_{oi} - \frac{1}{2} a_{oj} n^j n_i + \frac{1}{4} n_i a_{\mu}^{\mu}$$

harmonic gauge condition for  $\bar{h}_{\mu\nu}$ :  $\bar{h}_{\mu\nu} n^{\nu} = 0 \Rightarrow a_{\mu\nu} n^{\nu} = 0$

$$\Rightarrow n^0 a_{\mu 0} + a_{\mu j} n^j = 0 \Rightarrow a_{\mu 0} = -a_{\mu j} n^j$$

$$\mu=0: a_{00} = -a_{oj} n^j = a_{ij} n^i n^j$$

$$\text{and } a_{\mu}^{\mu} = -a_{00} + a_{ij} \delta^{ij} = (\delta^{ij} - a_{ij} n^i n^j) a_{ij} = \Lambda^{ij} a_{ij}$$

then:

$$b_0 = \frac{1}{2} a_{kl} n^k n^l + \frac{1}{4} \Lambda^{kl} a_{kl}$$

$$b_i = -a_{ij} n^j + \frac{1}{2} a_{kjl} n^k n^j n_i + \frac{1}{4} n_i \Lambda^{kl} a_{kl}$$

insert in eq. (19):

$$\begin{aligned} a_{ij}^{\pi\pi} &= a_{ij} + n_i (-a_{kj} n^k + \frac{1}{2} a_{kl} n^k n^l n_j + \frac{1}{4} n_j \Lambda^{kl} a_{kl}) \\ &\quad + n_j (-a_{ie} n^e + \frac{1}{2} a_{kle} n^k n^l n_i + \frac{1}{4} n_i \Lambda^{kl} a_{kl}) \\ &\quad - \delta_{ij} (n^0 b_0 + n^k b_k) \\ &= a_{ij} - a_{kj} n_i n^k - a_{ie} n_j n^e + a_{kle} n^k n^l n_i n_j + \frac{1}{2} n_i n_j \Lambda^{kl} a_{kl} - \delta_{ij} \frac{1}{2} \Lambda^{kl} a_{kl} \\ &= (\delta_{ij} - n_i n^k) (\delta_{jk} - n_j n^e) a_{ke} - \frac{1}{2} (\delta_{ij} - n_i n_j) \Lambda^{kl} a_{kl} \\ &= (\Lambda_i^k \Lambda_j^e - \frac{1}{2} \Lambda_{ij} \Lambda^{kl}) a_{ke} \stackrel{a_{ke}=a_{ek}}{\uparrow} \frac{1}{2} (\Lambda_i^k \Lambda_j^e + \Lambda_i^e \Lambda_j^k - \Lambda_{ij} \Lambda^{kl}) a_{kl} \end{aligned}$$

$\Lambda_{ij}^{kl} \Rightarrow (22)$

# 3.2 Geodesic Deviation

Ex P7

choose  $\tau^0 = 0$  &  $\tau^i$  is distance in lab. frame  
 for  $h_{\mu\nu} = 0$ : geodesics at rest stay at rest

$$\Lambda(U^\mu) = (1, 0, 0, 0) + \mathcal{O}(h)$$

in lab. frame:  $\tau = t, (U^\mu) = (1, 0, 0, 0), g_{\mu\nu} \approx \eta_{\mu\nu}, \Gamma^\mu_{\alpha\beta} \approx 0$

$$\frac{D^2 \tau^i}{D\tau^2} = \frac{d^2 \tau^i}{dt^2} = R^i_{\alpha\beta\gamma} U^\alpha U^\beta \tau^\gamma = R^i_{00j} \tau^j$$

Use (14):  $R^i_{00j} = \frac{1}{2} \partial_0^2 h^i_j + \mathcal{O}(h^2)$ , with  $h_{\mu\nu} = h^{\pi\mu}_{\nu\pi} = h^{\pi\mu}_{\nu\pi}, h^{\pi\mu}_{\nu\pi} = 0 = h^{\pi\mu}_{\nu\pi}$



$$\textcircled{1} = \frac{1}{2} (\partial_0^2 h^i_j + \partial_j^i \partial_j h^{\pi\pi}_{00} - \partial_j^i \partial_0 h^{\pi\pi}_{0j} - \partial_0 \partial_i h^{\pi\pi}_{0j})$$

on worldlines  
 $\Gamma^\mu_{\alpha\beta} = 0$   
 $\frac{d\tau^\mu}{d\tau} = 0$

$$\Rightarrow \frac{d^2 \tau^i}{dt^2} = \frac{1}{2} \tau^j \partial_0^2 h^i_j$$

Force field:  $F^i = \frac{1}{2} \partial_0^2 h^i_j \tau^j$

$\text{div } \vec{F} = \frac{\partial F^i}{\partial \tau^i} = \frac{1}{2} \partial_0^2 h^i_j \delta^j_i = 0$  & field lines have no source  
 i.e. no start and end



oscillation:  
 field lines "vanish"  
 to infinity and  
 come back.

in TT-gauge:

for  $h_{\mu\nu} = 0$ : geodesics at rest stay at rest  $\Lambda(U^\mu) = (1, 0, 0, 0) + \mathcal{O}(h)$

$\frac{d\tau^\mu}{d\tau} = 0 + \mathcal{O}(h)$ : no initial relative velocity

$$\Rightarrow \frac{D^2 \tau^i}{D\tau^2} = \frac{d^2 \tau^i}{dt^2} + \frac{d}{dt} (\Gamma^i_{\alpha\beta} U^\alpha \tau^\beta) + \Gamma^i \frac{d\tau}{d\tau} + \Gamma \Gamma$$

$$\tau = t + \mathcal{O}(h) \quad \frac{1}{2} (\partial_0^2 h^i_j + \partial_j^i \partial_j h^{\pi\pi}_{00} - \partial_j^i \partial_0 h^{\pi\pi}_{0j}) \cdot \tau^j + \mathcal{O}(h^2)$$

$$\approx \frac{d^2 \tau^i}{dt^2} + \frac{1}{2} \partial_0^2 h^i_j \tau^j = \frac{1}{2} \tau^j \partial_0^2 h^i_j$$

$$\Rightarrow \frac{d^2 \tau^i}{dt^2} \approx 0$$

coordinates move along  
 geodesics in TT-gauge!